

# Postpartum Depression: The most common medical complication of pregnancy/childbirth<sup>1-6</sup>



Postpartum depression, or PPD, is a **major depressive episode** that can occur during pregnancy or after giving birth, and may negatively impact maternal, birth and infant outcomes.<sup>7-9</sup>

PPD may have devastating consequences for the mother, and may negatively impact: **relationships with a partner | that partner's mental health | the infant's physical, mental and emotional development.**<sup>7, 8, 10-15</sup>

While the exact cause of PPD is unknown, a number of potential acute and chronic risk factors have been established, **including history of depression, hormonal fluctuations and chronic stress.**<sup>16-19</sup>



**In the U.S., estimates of new mothers identified with PPD each year vary by state from 8% to 20%, with an overall average of 11.5%.<sup>1</sup>**

Studies have shown that for low-income women and women of color, rates are significantly higher. Past history of depression or anxiety can increase the risk of PPD.<sup>20-22</sup>

## The symptoms & onset of PPD vary<sup>9, 22-24</sup>

### ONSET:

Expert opinions vary as to the timing of PPD onset. It is thought that PPD can begin during pregnancy or following childbirth, from within four weeks up to one year postpartum.

### COMMON SYMPTOMS INCLUDE:



Feeling sad, hopeless, empty, overwhelmed, worried or overly anxious



Avoiding friends or family



Losing interest in activities that are usually enjoyable



Having trouble bonding with your baby



Thoughts of harming yourself or your baby

## PPD often goes undiagnosed

Women receive medical care from a complex network of clinicians during the perinatal period, including primary care physicians, obstetrician-gynecologists (OB-GYNs), midwives, other specialists and advanced practice clinicians.<sup>25</sup>



**Yet, current estimates show that approximately half of PPD cases may go undiagnosed without screening.<sup>26-33</sup>**

Feelings of reluctance, shame, embarrassment, judgment or fear may prevent new mothers or their partners from seeking help. PPD is a medical condition, not a personal failure.<sup>34-37</sup>

Many leading medical organizations and government agencies recommend screening at least once during pregnancy and the postpartum period for all mothers.<sup>23,36-37</sup>

For more information, visit [www.SeePPD.com](http://www.SeePPD.com).

## References

1. Ko JY, Rockhill KM, Tong VT, et al. Trends in Postpartum Depressive Symptoms - 27 States, 2004, 2008, and 2012. *MMWR Morb Mortal Wkly Rep.* 2017;66:153-158.
2. DeSisto CL et al. Prevalence Estimates of Gestational Diabetes Mellitus in the United States, Pregnancy Risk Assessment Monitoring System (PRAMS), 2007-2010. *Prev Chronic Dis.* 2014;11:E10.
3. Centers for Disease Control and Prevention. Data on Selected Pregnancy Complication in the United States. 2017. Available at: <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-complications-data.htm>. Accessed May 8, 2017.
4. Centers for Disease Control and Prevention. Pregnancy-related mortality surveillance. 2017. Available at: [www.cdc.gov/reproductivehealth/MaternalInfantHealth/PMSS.html](http://www.cdc.gov/reproductivehealth/MaternalInfantHealth/PMSS.html).
5. Knight M et al. Trends in postpartum hemorrhage in high resource countries: a review and recommendations from the International Postpartum Hemorrhage Collaborative Group. *BMC Pregnancy Childbirth.* 2009;9:55.
6. Reddy UM et al. Serious maternal complications after early preterm delivery (24–33 weeks' gestation). *Am J Obstet Gynecol.* 2015;213:538. \*Data based on the 2-step screening process.
7. Grote NK, Bridge JA, Gavin AR, Melville JL, Iyengar S, Katon WJ. A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction. *Arch Gen Psychiatry.* 2010;67(10):1012-1024. doi:10.1001/archgenpsychiatry.2010.111.
8. Li D, Liu L, Odouli R. Presence of depressive symptoms during early pregnancy and the risk of preterm delivery: a prospective cohort study. *Hum Reprod.* 2009;24(1):146-153. doi:10.1093/humrep/den342.
9. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 5th ed. Washington, DC: American Psychiatric Association; 2013.
10. Fitelson E, Kim S, Baker AS, Leight K. Treatment of postpartum depression: clinical, psychological and pharmacological options. *Int J Womens Health.* 2010;3:1-14. doi:10.2147/IJWH.S6938.
11. Vismara L, Rollè L, Agostini F, et al. Perinatal Parenting Stress, Anxiety, and Depression Outcomes in First-Time Mothers and Fathers: A 3- to 6-Months Postpartum Follow-Up Study. *Frontiers in Psychology.* 2016;7. doi:10.3389/fpsyg.2016.00938.
12. Kerstis B, Engström G, Sundquist K, Widarsson M, Rosenblad A. The association between perceived relationship discord at childbirth and parental postpartum depressive symptoms: a comparison of mothers and fathers in Sweden. *Ups J Med Sci.* 2012;117(4):430-438. doi:10.3109/03009734.2012.684805.
13. Valla L, Wentzel-Larsen T, Smith L, Birkeland MS, Slinning K. Association between maternal postnatal depressive symptoms and infants' communication skills: A longitudinal study. *Infant Behavior and Development.* 2016;45:83-90.
14. Koutra K, Chatzi L, Bagkeris M, Vassilaki M, Bitsios P, Kogevinas M. Antenatal and postnatal maternal mental health as determinants of infant neurodevelopment at 18 months of age in a mother-child cohort (Rhea Study) in Crete, Greece. *Social Psychiatry and Psychiatric Epidemiology.* 2013;48(8):1335-1345. doi:10.1007/s00127-012-0636-0.
15. Goodman J.H. Paternal postpartum depression, its relationship to maternal postpartum depression, and implications for family health. *Journal of Advanced Nursing.* 2004. 45(1), 26-35.
16. Schiller CE, Meltzer-Brody S, Rubinow DR. The role of reproductive hormones in postpartum depression. *CNS Spectr.* 2015;20(1):48-59. doi:10.1017/S1092852914000480.
17. Yim IS, Tanner Stapleton LR, Guardino CM, Hahn-Holbrook J, Dunkel Schetter C. Biological and psychosocial predictors of postpartum depression: systematic review and call for integration. *Annu Rev Clin Psychol.* 2015;11:99-137. doi:10.1146/annurev-clinpsy-101414-020426.
18. Robertson E, Grace S, Wallington T, Stewart DE. Antenatal risk factors for postpartum depression: a synthesis of recent literature. *Gen Hosp Psychiatry.* 2004;26(4):289-295. doi:10.1016/j.genhosppsy.2004.02.006.
19. Silverman ME, Reichenberg A, Savitz DA, et al. The risk factors for postpartum depression: A population-based study. *Depress Anxiety.* 2017;34(2):178-187. doi:10.1002/da.22597.
20. Gress-Smith JL, Lueckel LJ, Lemery-Chalfant K, et al. Postpartum Depression Prevalence and Impact on Infant Health, Weight, and Sleep in Low-Income and Ethnic Minority Women and Infants. *Maternal and Child Health Journal.* 2012;16(4):887-893.
21. Chaudron LH, Szilagyi PG, Tang W, et al. Accuracy of Depression Screening Tools for Identifying Postpartum Depression Among Urban Mothers. *Pediatrics.* 2010;125(3):417-419.
22. Stewart DE, Robertson E, Dennis CL, et al. Postpartum depression: literature review of risk factors and interventions. University Health Network Women's Health Program; 2003. [http://www.who.int/mental\\_health/prevention/suicide/lit\\_review\\_postpartum\\_depression.pdf](http://www.who.int/mental_health/prevention/suicide/lit_review_postpartum_depression.pdf).
23. Screening for Perinatal Depression - ACOG. <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Screening-for-Perinatal-Depression>. Accessed May 3, 2018.
24. National Institute of Mental Health. Postpartum Depression Facts. <http://www.nimh.nih.gov/health/publications/postpartum-depression-facts/index.shtml>. Accessed April 6, 2018.
25. Kozhimannil KB, Fontaine P. Care From Family Physicians Reported by Pregnant Women in the United States. *Ann Fam Med.* 2013; 11(4):350-354.
26. Georgiopoulos AM. Routine Screening for Postpartum Depression. *J Fam Pract.* 2001;50:117-122.
27. Evins GG, et al. Postpartum depression: A comparison of screening and routine clinical evaluation. *Am J Obstet Gynecol.* 2000;182:1080-1082.
28. Cox EQ, Sowa NA, Meltzer-Brody SE, Gaynes BN. The Perinatal Depression Treatment Cascade: Baby Steps Toward Improving Outcomes. *J Clin Psychiatry.* 2016;77(9):1189-1200. doi:10.4088/JCP.15r10174.
29. Georgiopoulos AM, Bryan TL, Yawn BP, Houston MS, Rummans TA, Therneau TM. Population-based screening for postpartum depression. *Obstet Gynecol.* 1999;93(5 Pt 1):653-657.
30. Coates AO, Schaefer CA, Alexander JL. Detection of postpartum depression and anxiety in a large health plan. *J Behav Health Serv Res.* 2004;31(2):117-133.
31. Ko JY, Farr SL, Dietz PM, Robbins CL. Depression and treatment among US pregnant and nonpregnant women of reproductive age, 2005-2009. *J Womens Health (Larchmt).* 2012;21(8):830-836.
32. Spitzer RL, Williams JB, Kroenke K, Hornyak R, McMurray J. Validity and utility of the PRIME-MD Patient Health Questionnaire in assessment of 3000 obstetricgynecologic patients: the PRIME-MD Patient Health Questionnaire Obstetrics-Gynecology Study. *Am J Obstet Gynecol.* 2000;183(3):759-769.
33. Goodman JH, Tyer-Viola L. Detection, treatment, and referral of perinatal depression and anxiety by obstetrical providers. *J Womens Health (Larchmt).* 2010;19(3):477-490.
34. Byatt N, et al. Women's perspectives on postpartum depression screening in pediatric settings: a preliminary study. *Arch Womens Ment Health.* 2016;16:429-432.
35. National Institute for Health Care Management. Identifying and Treating Maternal Depression: Strategies & Considerations for Health Plans
36. Heneghan A.M., Morton S., DeLeone N.L. Paediatricians' attitudes about discussing maternal depression during a paediatric primary care visit. *Child: care, health and development.* 2006. 33, 3, 333-339.
37. Dunford E., Granger C. Maternal Guilt and Shame: Relationship to Postnatal Depression and Attitudes towards Help-Seeking. *Journal of Child and Family Studies.* 2017, Volume 26, Issue 6, pp 1692–1701.
38. Final Recommendation Statement: Depression in Adults: Screening - US Preventive Services Task Force. <https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/depression-in-adults-screening1>. Accessed May 3, 2018.
39. The Importance of Screening. <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Screening/Pages/The-Importance-of-Screening.aspx>. Accessed May 3, 2018.